



Enriched Xenon Observatory for double beta decay

D.Leonard, A.Piepke

Physics Dept, University of Alabama, Tuscaloosa AL

P.Vogel

Physics Dept Caltech, Pasadena CA

A.Bellerive, M.Bowcock, M.Dixit, I.Ekchtout, C.Hargrove, D.Sinclair, V.Strickland

Carleton University, Ottawa, Canada

W.Fairbank Jr., S.Jeng, K.Hall

Colorado State University, Fort Collins CO

M.Moe

Physics Dept UC Irvine, Irvine CA

D.Akimov, A.Burenkov, M.Danilov, A.Dolgolenko, A.Kovalenko, D.Kovalenko, G.Smirnov, V.Stekhanov

ITEP Moscow, Russia

J.Farine, D.Hallman, C.Virtue

Laurentian University, Canada

M.Hauger, F.Juget, L.Ounalli, D.Schenker, J-L.Vuilleumier, J-M.Vuilleumier, P.Weber

Physics Dept University of Neuchatel, Switzerland

M.Breidenbach, R.Conley, C.Hall, D.McKay, A.Odian, C.Prescott, P.Rowson, J.Sevilla, K.Skarpaas, K.Wamba

SLAC, Menlo Park CA


R.DeVoe, B.Flatt, G.Gratta, M.Green, F.LePort, R.Neilson, A.Pocar, S.Waldman, J.Wodin

Physics Dept Stanford University, Stanford CA

EXO will be large real time active source search for $\beta\beta$ decay of ^{136}Xe , using isotopically enriched Xenon.

Scientific goals are:

- 1) Measurement of yet unobserved $\beta\beta 2\nu$ decay of ^{136}Xe . Task: $T_{1/2} > 10^{22}$ y. **Particularly long.**
- 2) Test of the Heidelberg evidence for $\beta\beta 0\nu$ decay.
Expectation for ^{136}Xe [Ge $\pm 3\sigma$ range $(0.7-4.2) \cdot 10^{25}$ y]:
$$T_{1/2} = (0.57-3.4) \cdot 10^{25} \text{ y} \quad [\text{Rodin et al. PRC68 (2003) 044302 QRPA}]$$
$$= (0.66-4.0) \cdot 10^{25} \text{ y} \quad [\text{Staudt et al. EPL13 (1990) 31 QRPA}]$$
$$= (0.48-2.9) \cdot 10^{25} \text{ y} \quad [\text{Caurier et al. NPA654 (1999) 973c SM}]$$
- 3) Perform a search for $\beta\beta 0\nu$ decay sensitive enough to cover the allowed parameter space for an inverted mass hierarchy. $\langle m_{\beta\beta} \rangle < 10 \text{ meV} \rightarrow T_{1/2} > 4 \cdot 10^{28} \text{ y}$ [Rodin]
If (2) turns out a negative result.



EXO has chosen ^{136}Xe :

- **High Q-value: 2481 keV**
- **Active detection medium suited for charge collection plus high yield UV scintillator**
- **No need to grow crystals**
- **No long lived Xe isotopes to activate**
- **Can be easily transferred from one detector to another if new technologies become available**
- **Noble gas: easy(er) to purify re-purify during operation**
- **^{136}Xe has reasonable natural abundance: 8.9%**
- **^{136}Xe advantages for enrichment:**
 - noble gas (no chemistry involved)
 - centrifuge feed rate in gram/s, all mass useful
 - centrifuge efficiency $\sim \Delta m$. For Xe 4.7 amu
- **Ionization potentials Xe: 12.130 eV, Ba^+ : 5.212 eV, Ba^{++} : 10.004 eV $\rightarrow \beta\beta$ -decay product atom remains charged \rightarrow opens possibility of final state tagging**

Major challenges in $0\nu\beta\beta$ decay

1) Very large fiducial mass (tons)

need large-scale isotopic enrichment

2) Reduce and control backgrounds in qualitatively new ways

unlikely to gain big factors without new techniques

For a background scaling like Nt

$$\langle m_\nu \rangle \propto 1 / \sqrt{T_{1/2}^{0\nu\beta\beta}} \propto 1 / (Nt)^{1/4}$$

See M.Moe PRC44 (1991) 931

For no background


$$\langle m_\nu \rangle \propto 1 / \sqrt{T_{1/2}^{0\nu\beta\beta}} \propto 1 / \sqrt{Nt}$$

EXO plans to tag Ba^{++} ions born in $\beta\beta$ decays at vertex to operate in this regime.

Challenge: Background Control

“Real time” (Ionization and scintillation)


- + Energy/tracking information
- γ backgrounds
- Isotopic enrichment (usu.)
- Exposure (Nt) limited by time



Final state ID

- + Very specific signature
- + Large fiducial masses
- No decay-mode distinction

EXO is designed to combine these two techniques




See M.Moe PRC44 (1991) 931

**Leakage of $\beta\beta 2\nu$ -
events into $\beta\beta 0\nu$ -
analysis window.**

**Depends in 5.8th
power on σ .**


**Even with Ba
tagging important
background.**



EXO road map

- Build a 200 kg liquid Xe tracking TPC to demonstrate E resolution, low background, and tracking.
 - Largest liquid Xe detector ever build
 - One of largest TPCs
 - Largest $\beta\beta$ experiment by factor 20 compared to HD-MO
- EXO-200 will initially not be equipped with Ba tagging.
- Will be operated at moderate overburden at WIPP
[1585 mw.e. $4.8 \cdot 10^{-7} \mu/(\text{cm}^2 \text{ s})$ Esch et al. astro-ph/0408486] to measure $\beta\beta 2\nu$ decay and test Heidelberg $\beta\beta 0\nu$ evidence
- Planned installation at WIPP: end of 2006
- In parallel develop Ba final state tagging
- After proof of feasibility build a 1 to 10 ton enriched Xe detector to probe inverted mass hierarchy

The roadmap to the background free discovery of Majorana neutrinos and the neutrino mass scale




EXO-200 Progress


EXO-200 at WIPP

Modules are then moved to NExA and placed in position and connected. Distance traveled ~ 2100 ft.

Modules arrive here and are transferred to WIPP flatcar. Flatcar is moved to waste handling shaft and lowered into mine. Flatcar load rating: 67,000 lbs.



Energy resolution improvement in LXe



Ionization alone:
 $\sigma(E)/E = 3.8\%$ @ 570 keV
or 1.8% @ $Q_{\beta\beta}$

Ionization & Scintillation:
 $\sigma(E)/E = 3.0\%$ @ 570 keV
or 1.4% @ $Q_{\beta\beta}$
(a factor of 2 better than the Gotthard TPC)

E.Conti et al. Phys. Rev. B (68) 054201

**EXO-200 will collect 3-4 times
as much scintillation...
further improvement possible**

Massive effort on material radioactivity qualification

- **NAA^a**
- **Low background γ -spectroscopy^b**
- **α -counting^c**
- **Radon counting^d**
- **High performance GD-MS and ICP-MS^e**

**Th/U Sensitivity for
Teflon (TPC body):
<0.3 ppt or
1 and 4 $\mu\text{Bq/kg}$**

**Online database for collaborators at
present includes ~130 entries**

**MC simulation of backgrounds
Alabama & SLAC**

^a Alabama using MIT reactor




^b Neuchatel, Alabama


^c Alabama, Stanford, SLAC, Carleton

^d Laurentian


^e Commercial, Canadian Inst. Standards

Cryostat and gantry system views





EXO-200: a 200 kg LXe TPC with scintillation readout in a ultra-low background cryostat/shielding



Cryostat being fabricated at SDMS (Grenoble)

After machining and welding plates are returned to shielded storage





Modular Clean Rooms at Stanford



EXO-200 TPC Basics

- Measure both ionization and scintillation light to get best energy resolution
 - Info on event topology important for background separation.
 - Test ability to measure position of decay for Ba^+ extraction in full EXO
- The detector is a cylinder of 44 cm ID by 44 cm inner length.
- The cylinder is split by a cathode plane at the center so there are two symmetric drift regions. The cathode runs at negative HV.
 - Max HV is $\sim 70\text{kV}$ ($\sim 3.5\text{ kV/cm}$ drift). Energy resolution improves with drift field, but there are arguments that separation of 1 vs. 2 primary electrons might be better at lower fields.
 - field optimization is an important mission of EXO-200
- Readout “style”:
 - Crossed wires, $100\mu\text{m}$ wires, 3mm pitch, ganged in groups of 3 (48ch. x, 48ch. y), total 96 ch. per 1/2 detector
 - APDs for scintillation readout


Teflon chamber R&D



Metal-teflon seal

Recirculation line

Teflon chamber



Unmounted LAAPD from Advanced Photonix

Special gold-less production for EXO-200

APDs are ideal for our application:

- very clean & light-weight,
- very sensitive to VUV

$QE > 1$ at 175nm

Gain ~ 100-150


$V \sim 1500V$

$\Delta V < \pm 0.5V$



$\Delta T < \pm 1K$ APD is the driver for temperature stability

Leakage current OK cold

Channel count: 2×300




Mounted APD Layout




Ba⁺ Grabbing & Tagging R&D


Single atom spectroscopy at Stanford and Colorado State





Blue and red lasers
with diagnostics



Single-ion detection in presence of He background gas



Initial R&D on grabbing/release uses ^{230}U source and α -counting at SLAC



Ba grabbing and tagging R&D

- **Good quality tagging in presence of some Xe/He gas**
- **He stabilization of Xe-induced unloading**
- **High efficiency trap-loading scheme (linear trap)**
- **Linear trap construction close to final device**
- **Single Ba ion source development**

Ba tagging

- **Experience grabbing on Xe-ice and metal tips**
- **Field emission release from STM tip, further cryotip developments**

Ba grabbing

Done
In progress

In addition continue R&D on tagging in LXe:
very high rewards if successful.

EXO-200 Majorana mass sensitivity

Assumptions:

- 1) 200kg of Xe enriched to 80% in ^{136}Xe
- 2) $\sigma(E)/E = 1.4\%$ obtained in EXO R&D, Conti et al Phys Rev B 68 (2003) 054201
- 3) Low but finite radioactive background:
20 events/year in the $\pm 2\sigma$ interval centered around the 2.481MeV endpoint
- 5) Negligible background from $2\nu\beta\beta$ ($T_{1/2} > 1 \cdot 10^{22}\text{yr}$ R.Bernabei et al. measurement)

Case	Mass (ton)	Eff. (%)	Run Time (yr)	σ_E/E @ 2.5MeV (%)	Radioactive Background (events)	$T_{1/2}^{0\nu\beta\beta}$ (yr, 90%CL)	Majorana mass (eV)	
							QRPA	NSM
Prototype	0.2	70	2	1.6*	40	$6.4 \cdot 10^{25}$	0.27†	0.38♦

What if Klapdor's observation is correct ?

Central value $T_{1/2}(\text{Ge}) = 1.2^{+3}_{-0.5} \cdot 10^{25}$, $\pm 3\sigma$ range (0.24eV – 0.58eV)
(Phys. Lett. B 586 (2004) 198-212)

In 200kg EXO, 2yr:

• Worst case (QRPA, upper limit) 15 events on top of 40 events bkgd $\rightarrow 2\sigma$

• Best case (NSM, lower limit) 162 events with 40 bkgd $\rightarrow 8.5\sigma$

EXO Majorana mass sensitivity

Assumptions:

- 1) 80% enrichment in ^{136}Xe
- 2) Intrinsic low background + Ba^+ tagging eliminate all radioactive background
- 3) Energy resolution only used to separate the 0ν from 2ν modes:
Select 0ν events in a $\pm 2\sigma$ interval centered around the 2.481 MeV endpoint
- 4) Use for $2\nu\beta\beta$ $T_{1/2} > 1 \cdot 10^{22}$ yr (Bernabei et al. measurement)

Case	Mass (ton)	Eff. (%)	Run Time (yr)	σ_E/E @ 2.5MeV (%)	$2\nu\beta\beta$ Background (events)	$T_{1/2}^{0\nu\beta\beta}$ (yr, 90%CL)	Majorana mass (meV) QRPA [‡] NSM [#]	
Conservative	1	70	5	1.6*	0.5 (use 1)	2×10^{27}	50	68
Aggressive	10	70	10	1 [†]	0.7 (use 1)	4.1×10^{28}	11	15

* $\sigma(E)/E = 1.4\%$ obtained in EXO R&D, Conti et al Phys Rev B (68) (2003) 054201

[†] $\sigma(E)/E = 1.0\%$ considered as an aggressive but realistic guess with large light collection area

[‡] Rodin et al Phys Rev C 68 (2003) 044302

[#] Courier et al. Nucl Phys A 654 (1999) 973c